

Two-Component Formulation of the Wheeler-DeWitt Equation

Ali Mostafazadeh*

Theoretical Physics Institute, University of Alberta,
Edmonton, Alberta, Canada T6G 2J1.

Abstract

The Wheeler-DeWitt equation for the minimally coupled FRW-massive-scalar-field minisuperspace is written as a two-component Schrödinger equation with an explicitly ‘time’-dependent Hamiltonian. This reduces the solution of the Wheeler-DeWitt equation to the eigenvalue problem for a non-relativistic one-dimensional harmonic oscillator and an infinite series of trivial algebraic equations whose iterative solution is easily found. The solution of these equations yields a mode expansion of the solution of the original Wheeler-DeWitt equation. Further analysis of the mode expansion shows that in general the solutions of the Wheeler-DeWitt equation for this model are doubly graded, i.e., every solution is a superposition of two definite-parity solutions. Moreover, it is shown that the mode expansion of both even and odd-parity solutions is always infinite. It may be terminated artificially to construct approximate solutions. This is demonstrated by working out an explicit example which turns out to satisfy DeWitt’s boundary condition at initial singularity.

*E-mail: alimos@phys.ualberta.ca

1 Introduction

It is well-known that the Wheeler-DeWitt equation for the minimally coupled FRW-massive-scalar-field ¹ minisuperspace can be written in the form of the Klein-Gordon equation ² in $(1+1)$ dimensions [1],

$$\left[-\frac{\partial^2}{\partial \alpha^2} + \frac{\partial^2}{\partial \phi^2} + \kappa e^{4\alpha} - m^2 e^{6\alpha} \phi^2 \right] \psi(\alpha, \phi) = 0, \quad (1)$$

where $\alpha = \ln a$, a is the scale factor of the FRW metric

$$ds^2 = -dt^2 + a^2(t) \left[dr^2 + S_\kappa(r)(d\theta^2 + \sin^2 \theta d\varphi^2) \right],$$

$$S_\kappa := \begin{cases} \sinh r & \text{for } \kappa = -1 : \text{ open universe} \\ r & \text{for } \kappa = 0 : \text{ flat universe} \\ \sin r & \text{for } \kappa = 1 : \text{ closed universe,} \end{cases}$$

and ϕ is a real scalar field of mass m . It is also well-known that unlike the massless case [2] exact solutions of this equation are not known. The approximate solutions are usually obtained using the semiclassical (WKB) methods, [3].

Recently, I have used a two-component formulation of the Klein-Gordon equation to study the geometrical phases induced on a Klein-Gordon field due to a dynamical background spacetime geometry and electromagnetic or scalar interactions, [4]. In the present article, I shall try to apply the same approach for the treatment of the Wheeler-DeWitt equation (1).

2 Two-component Form of the Wheeler-DeWitt Equation

The idea of the two-component formalism [5] is to transform the Klein-Gordon equation which is of second order in time-derivative, into a system of two coupled differential equations which are first order in the time-derivative. The latter may be viewed as a Schrödinger equation. In this way one can apply the well-developed techniques of solving the Schrödinger equation to obtain solutions of the Klein-Gordon equation. The application of the two-component formalism for the Klein-Gordon equation is described in detail in Ref. [4]. Here I only include the results which are relevant for the solution of Eq. (1).

¹The scalar field is assumed to be homogeneous.

²This equation corresponds to simplest choice of factor ordering.

Eq. (1) can be expressed in the form:

$$\left[\eta^{\mu\nu} \frac{\partial}{\partial X^\mu} \frac{\partial}{\partial X^\nu} + V(X) \right] \psi = 0 , \quad (2)$$

where $\mu, \nu = 0, 1$, $(\eta^{\mu\nu}) := \text{diag}(-1, 1)$, $X^0 := \alpha$, $X^1 := \phi$, and

$$V(X) := \kappa e^{4X^0} - m^2 e^{6X^0} (X^1)^2 = \kappa e^{4\alpha} - m^2 e^{6\alpha} \phi^2 = \kappa a^4 - m^2 a^6 \phi^2 . \quad (3)$$

Hence, α plays the role of ‘time.’ Furthermore, defining

$$\Psi := \frac{1}{\sqrt{2}} \begin{pmatrix} \psi + i\dot{\psi} \\ \psi - i\dot{\psi} \end{pmatrix} , \quad (4)$$

one can easily check that Eq. (2) is equivalent to the Schrödinger equation

$$i\dot{\Psi} = H\Psi , \quad (5)$$

with the explicitly ‘time’-dependent Hamiltonian:

$$H = \frac{1}{2} \begin{pmatrix} 1 + D & -1 + D \\ 1 - D & -1 - D \end{pmatrix} , \quad (6)$$

where a dot represents differentiation with respect to $X^0 = \alpha$ and

$$D := -\frac{\partial^2}{\partial (X^1)^2} - V = -\frac{\partial^2}{\partial \phi^2} - V . \quad (7)$$

The two-component Hamiltonian (6) has the following remarkable properties:

- It belongs to the (dynamical) Lie algebra $su_s(1, 1) \otimes su_u(1, 1)$. Here $su_s(1, 1)$ is the standard representation of $su(1, 1)$ with the basis

$$K_3 = J_3 = \frac{1}{2} \sigma_3 , \quad K_a = iJ_a = \frac{i}{2} \sigma_a , \quad (a = 1, 2) ,$$

where σ_i are Pauli matrices, and $su_u(1, 1)$ denotes the unitary representation of $su(1, 1)$ constructed out of the position x and momentum p operators of the one-dimensional quantum mechanics, with the basis $\{(x^2 - p^2)/4, (x^2 + p^2)/4, (xp + px)/4\}$, [6]. This is easily seen, by writing H in the form $H = K_- + K_+ D$, where $K_\pm := K_3 \pm K_2$, and observing that the operator D belongs to $su_u(1, 1)$.

— It is the square root of the operator D , in the sense that it satisfies

$$H^2 = \begin{pmatrix} D & 0 \\ 0 & D \end{pmatrix} = D. \quad (8)$$

This is reminiscent of Dirac's treatment of the non-interacting Klein-Gordon equation in $(3+1)$ -dimensional Minkowski space, where the Dirac operator plays the role of the square root of the Klein-Gordon operator.

Next let us examine the eigenvalues E_n and eigenvectors Ψ_n of the Hamiltonian H . It is not difficult to show that due to the particular form of H , the eigenvalue equation

$$H\Psi_n = E_n\Psi_n, \quad (9)$$

leads to the following relations:

$$\Psi_n = \frac{1}{\sqrt{2}} \begin{pmatrix} 1 + E_n \\ 1 - E_n \end{pmatrix} \Phi_n, \quad D\Phi_n = E_n^2\Phi_n. \quad (10)$$

Therefore the eigenvalue problem for H is reduced to that of D . Substituting Eq. (7) in the second equation in (10), one has:

$$\left[-\frac{1}{2} \frac{\partial^2}{\partial \phi^2} + \frac{1}{2} \omega^2 \phi^2 \right] \Phi_n = \mathcal{E}_n \Phi_n, \quad (11)$$

where $\omega := me^{3\alpha} = ma^3$ and $\mathcal{E}_n := (E_n^2 + \kappa e^{4\alpha})/2 = (E_n^2 + \kappa a^4)/2$. Eq. (11) is the eigenvalue equation for a simple harmonic oscillator with frequency ω and unit mass. Hence:

$$\mathcal{E}_n = \omega \left(n + \frac{1}{2} \right) = ma^3 \left(n + \frac{1}{2} \right), \quad (12)$$

$$\Phi_n = N_n H_n(\sqrt{\omega}\phi) e^{-\frac{1}{2}\omega\phi^2} = N_n H_n(\sqrt{ma^3}\phi) e^{-\frac{1}{2}ma^3\phi^2}, \quad (13)$$

where H_n are Hermite polynomials and N_n are normalization constants. The latter are determined by the requirement that Φ_n be normalized with the usual $L^2(\mathbb{R})$ norm. This leads to

$$N_n = \left[\frac{\omega}{\pi(2^n n!)^2} \right]^{1/4} = \left[\frac{ma^3}{\pi(2^n n!)^2} \right]^{1/4}.$$

Note that Φ_n and consequently Ψ_n tend to zero as $a \rightarrow 0$ or $a \rightarrow \infty$.

Substituting Eq. (12) in the expression for $\mathcal{E}_n$, one has:

$$E_{\pm n} = \pm \sqrt{\omega(2n+1) - \kappa e^{4\alpha}} = \pm \sqrt{me^{3\alpha}(2n+1) - \kappa e^{4\alpha}} = \pm a \sqrt{ma(2n+1) - \kappa a^2}. \quad (14)$$

Thus, for $\kappa = -1$ or 0 , $E_{\pm n}$ are real, whereas for $\kappa = 1$ and $a > m$, there are eigenvectors $\Psi_{\pm n}$ with imaginary eigenvalue. They correspond to the integers n satisfying $m(2n+1) < a$.

So far, I have given the solution for the eigenvalue problem for the Hamiltonian H . The eigenvectors of H , however, do not solve the Schrödinger equation. This is mainly because H is explicitly ‘time’-dependent. The general solution of the Schrödinger equation for an arbitrary explicitly time-dependent Hamiltonian is not known. However, in the cases where the eigenvalue problem for the Hamiltonian is solvable, there are techniques based on the adiabatic approximation [7, 8, 9] which may be used to obtain approximate and occasionally exact solutions. These techniques can only be employed if the Hamiltonian is self-adjoint. In order to examine the self-adjointness of the Hamiltonian H , one first needs a notion of an inner product on the space $\mathcal{H}$ of two-component wave functions Ψ .

3 Hilbert Space Structure of $\mathcal{H}$

As a vector space, the space $\mathcal{H}$ of two-component wave functions is isomorphic to $L^2(\mathbb{R}) \oplus L^2(\mathbb{R})$. Here by the vector space $L^2(\mathbb{R}) \oplus L^2(\mathbb{R})$, I mean the vector space inner sum of two copies of $L^2(\mathbb{R})$, not the Hilbert space inner sum of these spaces where the sum is also endowed with a canonical inner product. Furthermore, in what follows I allow for negative norms and the term Hilbert space refers to a complete not necessarily positive-definite inner product space [10]. $\mathcal{H} = L^2(\mathbb{R}) \oplus L^2(\mathbb{R})$ may be viewed as a restriction on the solutions of the Wheeler-DeWitt equation, since for finite α the elements of $L^2(\mathbb{R}) \oplus L^2(\mathbb{R})$ vanish at ‘spatial’ infinity, i.e., $\phi = \pm\infty$. This choice seems, however, to be a natural and at the same time a rather general one. It is only restrictive as far as the boundary condition at ‘spatial’ infinity is concerned. It is still quite general, for it does not correspond to any fixed boundary conditions at $\alpha = \pm\infty$. Furthermore, although $\mathcal{H}$ has two vector subspaces with L^2 inner products, an inner product on $\mathcal{H}$ is not a priori fixed.

A Hermitian inner product on $\mathcal{H}$ may be given by the inner product on $L^2(\mathbb{R})$ and a Hermitian two-by-two matrix h . A particularly advantageous choice for h is $h = \sigma^3 = \text{diag}(1, -1)$, i.e.,

$$(\Psi, \tilde{\Psi}) := \langle \Psi^1 | \tilde{\Psi}^1 \rangle - \langle \Psi^2 | \tilde{\Psi}^2 \rangle, \quad (15)$$

where $\langle | \rangle$ denotes the inner product on $L^2(\mathbb{R})$. This choice, on the one hand, renders H self-

adjoint. On the other hand, it reduces to the Klein-Gordon inner product³ for the solutions (4) of the Schrödinger equation (5).

Adopting this inner product on $\mathcal{H}$, one can compute

$$(\Psi_{\pm m}, \Psi_{\pm n}) = (E_{\pm m}^* + E_{\pm n})\delta_{mn} . \quad (16)$$

Here I have used (10) and the fact that Φ_n are orthonormal. In particular, $(\Psi_{\pm n}, \Psi_{\pm n}) = 2\text{Re}(E_{\pm n}) = \pm 2|\text{Re}(E_n)|$. Hence, the eigenvectors $\Psi_{\pm n}$ with imaginary eigenvalues (if they exist) have zero norm; they are null vectors. This is of course to be expected since the Hamiltonian is self-adjoint.⁴ Furthermore, note that the null eigenvectors are only present for the $\kappa = 1$ case.

Another interesting consequence of Eq. (16) is that

$$(\Psi_{\pm n}, \Psi_{\mp n}) = \begin{cases} 0 & \text{for } E_n \text{ real} \\ E_{\mp n} & \text{for } E_n \text{ imaginary.} \end{cases} \quad (17)$$

Hence, if we view $\mathcal{H}$ as the linear span of $\Psi_{\pm n}$, it can be orthogonally decomposed as

$$\mathcal{H} = \mathcal{H}^{(-)} \oplus \mathcal{H}^{(0)} \oplus \mathcal{H}^{(+)} , \quad (18)$$

where $\mathcal{H}^{(-)}$, $\mathcal{H}^{(0)}$, and $\mathcal{H}^{(+)}$ are the linear span of the negative-norm, null, and positive-norm eigenvectors. Note that this decomposition is ‘time’-dependent (it depends on $a = e^\alpha$), and not every null vector belongs to $\mathcal{H}^{(0)}$. Furthermore, Eq. (17) implies that the basis $\{\Psi_n\}$ (resp. $\{\Psi_{-n}\}$) of $\mathcal{H}^{(+)}$ (resp. $\mathcal{H}^{(-)}$) is orthogonal, whereas $\mathcal{H}_0$ can be orthogonally decomposed into the sum of two-dimensional subspaces $\mathcal{H}_n^{(0)} = \text{Span}(\Psi_n \oplus \Psi_{-n})$ with $n \leq (a/m - 1)/2$, namely

$$\mathcal{H}^{(0)} = \bigoplus_{n=1}^N \mathcal{H}_n^{(0)} , \quad (19)$$

where $N := [(a/m - 1)/2]$ is the largest integer part of $(a/m - 1)/2$.

4 Adiabatic Approximation

Consider the ansatz

$$\Psi = e^{i\zeta\mathcal{N}} \Psi_{\mathcal{N}} , \quad (20)$$

³For more details see Ref. [4].

⁴Note that it is the non-positive definiteness of the corresponding norm that allows for imaginary eigenvalues and null eigenvectors to exist.

for the solution of the Schrödinger equation (5), where $\zeta_{\mathcal{N}}$ is a complex-valued function of α , and $\mathcal{N} = \pm n \in \mathbb{Z}$. Plugging (20) in (5) and making use of the eigenvalue equation (9) and the form (10) of the eigenvectors $\Psi_{\mathcal{N}}$, one has:

$$\begin{aligned} \left[-(\dot{\zeta}_{\mathcal{N}} + E_{\mathcal{N}})(1 + E_{\mathcal{N}}) + i\dot{E}_{\mathcal{N}} \right] \Phi_n + i(1 + E_{\mathcal{N}})\dot{\Phi}_n &= 0 , \\ \left[-(\dot{\zeta}_{\mathcal{N}} + E_{\mathcal{N}})(1 - E_{\mathcal{N}}) - i\dot{E}_{\mathcal{N}} \right] \Phi_n + i(1 - E_{\mathcal{N}})\dot{\Phi}_n &= 0 , \end{aligned}$$

Adding and subtracting both sides of these equations, one finds:

$$(\dot{\zeta}_{\mathcal{N}} + E_{\mathcal{N}})\Phi_n - i\dot{\Phi}_n = 0 , \quad \dot{E}_{\mathcal{N}} = 0 . \quad (21)$$

Here the first equation implies that for $\tilde{n} \neq n$, $A_{\tilde{n}n} := i\langle \Phi_{\tilde{n}} | \dot{\Phi}_n \rangle = 0$. This requirement together with the second relation in (21) signify the conditions under which the adiabatic approximation would yield an exact solution of the Schrödinger equation. Clearly, $\dot{E}_{\mathcal{N}} \neq 0$ for the case under consideration. Thus the adiabatic approximation cannot be exact. However, approximate solutions may be sought if $\dot{E}_{\mathcal{N}}$ and $A_{\tilde{n}n}$ are small.

Performing the necessary calculations, one can show that for arbitrary non-negative integers $\tilde{n}$ and n :

$$A_{\tilde{n}n} := i\langle \Phi_{\tilde{n}} | \dot{\Phi}_n \rangle = \frac{3i}{4} \left[\sqrt{n(n-1)} \delta_{\tilde{n},n-2} - \sqrt{(\tilde{n})(\tilde{n}-1)} \delta_{\tilde{n},n+2} \right] , \quad (22)$$

$$\dot{E}_{\mathcal{N}} = \pm \frac{3 \left[n + \frac{1}{2} - \frac{2\kappa a}{3m} \right] \sqrt{m} a^{3/2}}{\sqrt{2n+1 - \frac{\kappa a}{m}}} , \quad \text{where } \mathcal{N} = \pm n . \quad (23)$$

Eq. (22) indicates that the adiabatic approximation is not valid, i.e., the solutions of the form (20) do not exist.

Another consequence of Eq. (22) is that in a mode expansion of a solution Ψ of Eq. (5), the even modes $\Psi_{2\mathcal{N}}$ and the odd modes $\Psi_{2\mathcal{N}+1}$ do not mix in the course of evolution. This suggests the existence of a double grading of the set of solutions. In particular one has the *even* and *odd* solutions:

$$\Psi_{\text{even}} = \sum_{\mathcal{N}=-\infty}^{\infty} Z_{2\mathcal{N}} \Psi_{2\mathcal{N}} , \quad \Psi_{\text{odd}} = \sum_{\mathcal{N}=-\infty}^{\infty} Z_{2\mathcal{N}+1} \Psi_{2\mathcal{N}+1} . \quad (24)$$

5 Mode Expansion of the Solutions

Consider expressing the solution Ψ of the two-component Wheeler-DeWitt equation (5) as a linear combination of the eigenfunctions $\Psi_{\mathcal{N}}$ of the Hamiltonian:

$$\Psi = \sum_{\mathcal{N}=-\infty}^{\infty} Z_{\mathcal{N}} \Psi_{\mathcal{N}}, \quad Z_{\mathcal{N}} = Z_{\mathcal{N}}(\alpha). \quad (25)$$

Substituting this equation in (5) and using Eqs. (9) and (10) to simplify the result, one finds

$$\sum_{n=0}^{\infty} [(\dot{\sigma}_n + i\sigma_{-n}E_n)\Phi_n + \sigma_n\dot{\Phi}_n] = 0, \quad (26)$$

$$\sum_{n=0}^{\infty} [(\dot{\sigma}_{-n}E_n + i\sigma_nE_n^2 + \sigma_{-n}\dot{E}_n)\Phi_n + \sigma_{-n}E_n\dot{\Phi}_n] = 0, \quad (27)$$

where

$$\sigma_{\pm n} = \sigma_{\pm n}(\alpha) := Z_n \pm Z_{-n}, \quad n = 0, 1, 2, \dots$$

Note that in general $Z_{-0} \neq Z_0$ and $\sigma_{-0} \neq \sigma_0$.

Next take the inner product of both sides of Eqs. (26) and (27) with $\Phi_{\tilde{n}}$. Making use of Eq. (22) and orthonormality of Φ_n , one has:

$$\dot{\sigma}_n + iE_n\sigma_{-n} + \frac{3}{4}\sqrt{(n+1)(n+2)}\sigma_{n+2} - \frac{3}{4}\sqrt{n(n-1)}\sigma_{n-2} = 0, \quad (28)$$

$$E_n\dot{\sigma}_{-n} + iE_n^2\sigma_n + \dot{E}_n\sigma_{-n} + \frac{3}{4}\sqrt{(n+1)(n+2)}E_{n+2}\sigma_{-(n+2)} - \frac{3}{4}\sqrt{n(n-1)}E_{n-2}\sigma_{-(n-2)} = 0, \quad (29)$$

where $\sigma_{-(0)} = \sigma_{-0}$.

In order to analyze these equations further, let us solve for σ_{-n} from (28) and substitute the result in (29). Doing the same for $\sigma_{-(n\pm 2)}$, one obtains:

$$\sigma_{-n} = \frac{i}{E_n} \left[\dot{\sigma}_n + \frac{3}{4}\sqrt{(n+1)(n+2)}\sigma_{n+2} - \frac{3}{4}\sqrt{n(n-1)}\sigma_{n-2} \right], \quad (30)$$

$$\begin{aligned} \ddot{\sigma}_n + [E_n^2 - \frac{3}{2}(n^2 + n + 1)]\sigma_n + \frac{3}{2}\sqrt{(n+1)(n+2)}\dot{\sigma}_{n+2} \\ - \frac{3}{2}\sqrt{n(n-1)}\dot{\sigma}_{n-2} + (\frac{3}{4})^2\sqrt{(n+1)(n+2)(n+3)(n+4)}\sigma_{n+4} + \\ (\frac{3}{4})^2\sqrt{n(n-1)(n-2)(n-3)}\sigma_{n-4} = 0. \end{aligned} \quad (31)$$

Note that Eq. (31) links only σ_{2n} 's, with $n = 0, 1, \dots$, thus making the analysis of the coefficients of non-negative modes independent of those of the negative modes.

Eqs. (30) and (31) clearly display the splitting of the solutions of the Wheeler-DeWitt equation into even and odd solutions. Since these equations are linear and there is no correlation between the even and odd solutions, every solution is a linear combination of an even and an odd solution. Hence without loss of generality, one can restrict to the analysis of Ψ_{even} and Ψ_{odd} .

Before engaging in this analysis, however, I would like to note that the mode expansion (25) for the two-component wave function Ψ , translates into

$$\psi = \sum_{n=0}^{\infty} \sigma_n \Phi_n, \quad (32)$$

for the one-component wave function. Thus, it is the coefficients of the zero and positive modes that determine the solution of the Wheeler-DeWitt equation (1). The negative modes are related to the derivative of ψ with respect to α . Hence, they are also uniquely determined once the zero and positive modes are given.

In the following sections, I shall show that the mode expansion of the solution of the Wheeler-DeWitt equation (1) is always infinite. This is another way of demonstrating the fact that the adiabatic approximation, which corresponds to including only a single mode in this expansion, always fails. Nevertheless, if one tries to blindly apply the adiabatic approximation or a slightly improved version of it in which one sets all $\sigma_{\pm n}$ to zero, except $\sigma_{\pm n_*}$ for some n_* , then Eqs. (28) and (29) yield $\ddot{\sigma}_{n_*} + E_n^2 \sigma_{n_*} = 0$. This corresponds to the eigenvalue equation for a one-dimensional quantum particle of mass 1/2 and potential $\mathcal{V}(\alpha) := -E_n^2 = \kappa e^{4\alpha} - (2n+1)me^{3\alpha}$. Specifically, σ_{n_*} is the zero-energy eigenfunction. For $\kappa = 1$, $\mathcal{V}$ vanishes at $\alpha = -\infty$. It decreases to attain a negative minimum as α increases, and then monotonically increases to $+\infty$ as $\alpha \rightarrow +\infty$. Hence, if one requires the zero-energy eigenfunction to vanish at $\alpha = +\infty$ where $\mathcal{V} = +\infty$, then the mass m must satisfy a peculiar quantization condition. This is analogous to the mass quantization advocated in Ref. [11]. However, note that the basic hypothesis of this argument, namely the validity of the adiabatic approximation or alternatively the presence of only a single mode in the mode expansion (32), is never satisfied. Hence, the argument leading to the quantization of mass does not apply.

6 Even Solutions

Let us first prove:

Lemma 1: *An even solution $\Psi_{\text{even}} = \sum_{\mathcal{N}} Z_{2\mathcal{N}} \Psi_{2\mathcal{N}}$ is nontrivial, i.e., $\Psi_{\text{even}} \neq 0$, if and only if $\sigma_{\pm 0} \neq 0$.*

In order to prove this statement, one sets $n = 0$ in Eqs. (28) and (29). This leads to

$$\dot{\sigma}_0 + iE_0\sigma_{-0} + \frac{3\sqrt{2}}{4}\sigma_2 = 0, \quad (33)$$

$$E_0\dot{\sigma}_{-0} + \dot{E}_0\sigma_{-0} + iE_0^2\sigma_0 + \frac{3\sqrt{2}}{4}E_2\sigma_{-2} = 0. \quad (34)$$

Therefore, if $\sigma_{\pm 0} = 0$, both σ_2 and σ_{-2} must vanish. However, by virtue of Eqs. (31) and (30), this is sufficient to conclude that $\sigma_{2\mathcal{N}} = 0$, for all $\mathcal{N} \in \mathbb{Z}$. This completes the proof of Lemma 1.

Furthermore, one can prove

Lemma 2: *Every nontrivial even solution has an infinite mode expansion.*

The even solutions may nevertheless be constructed inductively in terms of $\sigma_{\pm 0}$. This is done by noting that Eqs. (30) and (31) yield trivial algebraic equations for $\sigma_{\pm 2(n+1)}$ in terms of $\sigma_{\pm 2n}$. Performing an induction on n , one can express the coefficients of all the modes as linear combinations of $\sigma_{\pm 0}$ and its derivatives. For example, one has

$$\sigma_2 = -\frac{1}{\sqrt{2}}\left(\frac{4}{3}\right)(\dot{\sigma}_0 + iE_0\sigma_{-0}), \quad (35)$$

$$\sigma_4 = \frac{1}{\sqrt{4!}}\left(\frac{4}{3}\right)^2 \left[\ddot{\sigma}_0 + 2iE_0\dot{\sigma}_{-0} + \left(\frac{3}{2} - E_0^2\right)\sigma_0 + 2i\dot{E}_0\sigma_{-0} \right]. \quad (36)$$

It is easily seen that as a result of this construction σ_{2n} involves the first n derivatives of σ_0 , first $n - 1$ derivatives of σ_{-0} , and the first $n - 1 - j$ derivatives of E_{2j} , for $j = 0, 1, \dots, n - 1$.

Since according to Eq. (23), $\dot{E}_n$ diverges at the zero of E_n , demanding regularity of σ_n would impose certain conditions on $\sigma_{\pm 0}$. In fact, in view of the appearance of E_n^2 in Eq. (31), which is analytic for every α , one can easily show that the only source of possible singularity is the term proportional to E_0 in the expression for σ_2 . Thus, in order to ensure the regularity of σ_{2n} , one must require that at the zero of E_0 , i.e., $\alpha = \ln(m/\kappa)$, $\sigma_{-0}E_0$ and all its derivatives must vanish.

In other words there must be an open neighborhood $\mathcal{O}$ of $\alpha = \ln(m/\kappa)$, in which $\sigma_{-0}(\alpha)$ is of the form $E_0(\alpha)f(\alpha)$ for some smooth (C^∞) function f . In particular $\sigma_{-0}(\ln(m/\kappa))$ must vanish. Note that for the flat ($\kappa = 0$) and open ($\kappa = -1$) cases, this condition is void and σ_{2n} are regular if and only if $\sigma_{\pm 0}$ are regular.

Finally note that termination of the mode expansion for a nontrivial even solution is equivalent to vanishing of at least three consecutive σ_{2n} 's. This would impose three functional conditions on $\sigma_{\pm 0}$ which are in general incompatible. This proves Lemma 2.

7 Odd Solutions

The structure of the odd solutions is similar to that of the even solutions. The role of $\sigma_{\pm 0}$ is played by $\sigma_{\pm 1}$. In particular, one has

Lemma 3: *An odd solution $\Psi_{\text{odd}} = \sum_{\mathcal{N}} Z_{2\mathcal{N}+1} \Psi_{2\mathcal{N}+1}$ is nontrivial, if and only if $\sigma_{\pm 1} := Z_1 \pm Z_{-1} \neq 0$.*

This is a direct consequence of Eqs. (28) and (31) which also provide an inductive construction of the mode coefficients $\sigma_{\pm(2n+1)}$ in terms of $\sigma_{\pm 1}$. For example one has:

$$\sigma_3 = -\frac{1}{\sqrt{3!}}\left(\frac{4}{3}\right)(\dot{\sigma}_1 + iE_1\sigma_{-1}), \quad (37)$$

$$\sigma_5 = \frac{1}{\sqrt{5!}}\left(\frac{4}{3}\right)^2 \left[\ddot{\sigma}_1 + 2iE_1\dot{\sigma}_{-1} + \left(\frac{9}{2} - E_1^2\right)\sigma_1 + 2i\dot{E}_1\sigma_{-1} \right]. \quad (38)$$

A natural consequence of this inductive construction is

Lemma 4: *Every nontrivial odd solution has an infinite mode expansion.*

The regularity condition on the odd solutions is analogous to the even case. One can again show that the only source of possible singularity is the term $E_1\sigma_{-1}$ in the expression for σ_3 . Hence the regularity of σ_{2n+1} and therefore ψ is ensured by requiring that in the vicinity of the zero of E_1 , namely $\alpha = \ln(3m/\kappa)$, σ_{-1} must be of the form $E_1(\alpha)g(\alpha)$, for some smooth function g .

The occurrence of two free functions of α in the construction of even and odd solutions is in complete agreement with the general structure of the Wheeler-DeWitt/Klein-Gordon equation (1) whose solutions depend on two boundary or two initial conditions. The latter are given by two

functions of α which in the above procedure can be used to determine $\sigma_{\pm 0}$ or $\sigma_{\pm 1}$ depending on whether the solution is even or odd. In general the solution will be the sum of an even and an odd part. In this case the boundary (initial) conditions would determine both $\sigma_{\pm 0}$ and $\sigma_{\pm 1}$. Once these functions are determined the above iterative construction yields the mode expansion of the general solution.

8 Real Solutions

Probably the best-known proposal for the solution of the Wheeler-DeWitt equation is that of Hartle and Hawking [12] who suggested the path integral expression $\psi \propto \int \exp(-I)$ for the ground state wave function. Here I stands for the Euclidean action. Hartle-Hawking wave function is by construction real.

It is not difficult to see that the reality of the one-component solution of the Wheeler-DeWitt equation implies the reality of the coefficient functions σ_n in its mode expansion (32). This is because Φ_n are already real. In view of this observation and Eq. (28), $E_n \sigma_{-n}$ must be imaginary. However, as discussed in section 2, for the open and flat universes ($\kappa = -1, 0$), E_n is always real so that σ_{-n} must be imaginary. For the close universe ($\kappa = 1$) E_n is real for $a \leq m(2n + 1)$ and imaginary otherwise. Hence in this case ($\kappa = 1$) the opposite holds for σ_{-n} , i.e., it is imaginary for $a < m(2n + 1)$ and real for $a > m(2n + 1)$.

These conditions can also be stated as the reality condition on σ_0 , σ_1 , σ_2 , and σ_3 which also generate the solutions. Therefore, they are rather trivial.

9 Approximate Solutions

As shown in the preceding sections, the mode expansion of the solutions never terminates. Therefore, one cannot use it to construct an exact solution unless one can actually find the generic expression for the mode coefficients and sum the series. This is an almost impossible task. However, one can terminate the mode expansion artificially and obtain an infinite class of approximate solutions, $\psi \approx \psi_{n*} := \sum_{n=0}^{n*} \sigma_n \Phi_n$. The domain of reliability of this solutions will in general depend on the boundary (initial) conditions, alternatively the functions $\sigma_{\pm 0}$ and $\sigma_{\pm 1}$.

In this section, I shall examine the simple choice $\sigma_{-0} = \sigma_{\pm 1} = 0$ and $\sigma_0 = 1$. Clearly, this corresponds to a real even solution whose mode coefficients can be obtained using Eq. (31). For example for the closed FRW case where $\kappa = 1$, one has

$$\begin{aligned}\sigma_2 &= 0, \quad \sigma_4 = \left(\frac{2}{3}\right)^{5/2} \left(\frac{3}{2} + a^4 - ma^3\right), \quad \sigma_6 = \left(\frac{32a^3}{81\sqrt{5}}\right)(3m - 4a), \\ \sigma_8 &= \frac{4}{243} \sqrt{\frac{2}{35}} (4a^8 - 40ma^7 + 36m^2a^6 + 324a^4 - 288ma^3 + 189), \\ \sigma_{10} &= -\frac{16}{10935\sqrt{7}} (96a^5 - 1000ma^4 + 744m^2a^3 + 3604a - 2655m)a^3.\end{aligned}$$

Substituting these relations in the expression for ψ_n and making use of the identity [13]

$$H_n(x) = 2^n x^n + \sum_{i=1}^{[n/2]} (-1)^i 2^{n-i} (2i-1)!! \binom{n}{2i} x^{n-2i},$$

to calculate the mode functions Φ_n , one can compute $\psi_0 = \psi_2, \psi_4, \psi_6, \psi_8$ and ψ_{10} . Figures 1 to 5 illustrate three-dimensional plots of these functions. Figures 6 to 9 are plots of the restriction of ψ_6, ψ_8 and ψ_{10} to the lines $a = 0.1, 1, 10, 100$. These figures show that for small a the finite mode approximation is quite reliable. In fact, as depicted in Fig. 6, for $a = 0.1$, the graphs of ψ_8 and ψ_{10} coincide. For larger a one may improve the approximation by including the contribution of higher modes.

Note also that initially ψ_n are non-zero for a wide range of values of ϕ . However, as a increases the width of this region, which is symmetric with respect to the line $\phi = 0$, shrinks. This is easily seen from Figures 6 to 9. Note also that for large a , the solutions picks about $\phi = 0$ and the approximate solutions become less reliable. As $a \rightarrow \infty$, all the approximate solutions seem to vanish except at $\phi = 0$.

10 Discussion and Conclusion

Probably the most important aspect of the two-component formulation of the Wheeler-DeWitt equation is the fact that the construction of the solutions in this formulation does not, a priori, require fixing a particular inner product or boundary conditions. The freedom in the choice of the inner product is reflected as the independence of the mode expansion from the choice of the Hermitian matrix h which together with the L^2 inner product on $L^2(\mathbb{R})$ determines the inner

product on $\mathcal{H} = L^2(\mathbb{R}) \oplus L^2(\mathbb{R})$. As far as the boundary conditions are concerned, besides the implicit restriction of vanishing boundary condition at ‘spatial’ infinity, the analysis presented here applies generally.

An interesting consequence of the application of the two-component formalism is a specific double grading of the solutions of the Wheeler-DeWitt equation. This allows one to write down the most general solution of this equation as a sum of definite-parity (even and odd) solutions whose structure can be easily understood. The two definite-parity parts of the solution do not mix during the evolution. In particular, if the initial wave function has definite parity, its parity remains intact throughout the evolution.

Unfortunately, the mode expansion of the solutions never terminates. Thus, a closed expression for an exact solution cannot be found in this way. However, approximate solutions can be constructed by artificially terminating the mode expansion. The domain of applicability of such approximations depend generally on the boundary (initial) conditions.

An alternative approximation is the adiabatic approximation which is, however, not valid. The ansatz of the adiabatic approximation, i.e., $\psi \approx \sigma_n(\alpha)\Phi_n(\alpha, \phi)$, is reminiscent of the application of the Born-Oppenheimer approximation in quantum cosmology originally advocated by Banks [14]. In this approach one views the gravitational degrees of freedom as the ‘slow’ variables and the matter degrees of freedom as the ‘fast’ variables and considers solutions of the form $\psi = \psi_{\text{gravity}}(\alpha)\psi_{\text{matter}}(\alpha, \phi)$. The results of this article indicate that for $\psi_{\text{matter}} = \Phi_n$ such an approximation cannot be valid.

Another notable implication of the results of this article is regarding the implicit boundary condition at the initial singularity $a = 0$. If one assumes that the coefficient functions σ_n are regular at $a = 0$, then as a consequence of the appearance of $a^{3/4}$ in the expression for Φ_n , the one-component wave-function ψ vanishes at $a = 0$. This is much in the spirit of an argument of DeWitt [15]. Here it follows as a direct consequence of a regularity condition (which may or may not hold). As seen from Figures 1-5, the choice $\sigma_{-0} = \sigma_{+1} = 0$ and $\sigma_0 = 1$ leads to approximate solutions which are regular at $a = 0$. They clearly satisfy DeWitt’s boundary conditions.

A problem which is not addressed in this article is that of the convergence of the mode expansion in $\mathcal{H}$. This is a difficult problem and I have not been able to find the necessary and sufficient

conditions under which the mode expansion converges. In fact, a rigorous convergence analysis requires a choice of a positive-definite inner product (norm) on $\mathcal{H}$. However, as demonstrated in the preceding section, there are regions of the minisuperspace in which the contribution from the higher order modes is small. This may be viewed as an indication of possible pointwise asymptotic convergence of the mode expansion in this regions. This argument is however far from being satisfactory and further investigation of this point is necessary.

Finally, I wish to emphasize the following points:

- The fact that the two-component Hamiltonian (6) belongs to the dynamical Lie algebra $su_s(1, 1) \otimes su_u(1, 1)$ suggests a further investigation of the solution of the corresponding Schrödinger equation using the group theoretical methods such as those employed in solving the Schrödinger equation for time-dependent harmonic oscillators, [16]. Recall that the Hamiltonian for a time-dependent harmonic oscillator belongs to $su_u(1, 1)$, [6].
- The two-component approach to Wheeler-DeWitt equation has a wider domain of applicability than the simple minisuperspace model studied in this article. For example, the addition of an interaction potential for the scalar field would only complicate the defining equation for Φ_n and the inductive construction of the mode expansion. The general structure of the method would however remain intact. In particular, the ϕ -dependence of the solutions would still be given by Φ_n . More generally, for any other minisuperspace, in the gauge in which the shift vector corresponding to DeWitt's supermetric [15] vanishes, the two-component Hamiltonian is of the form (6), where D is some second order elliptic operator. Therefore, even in this case the developments reported in this article are essentially applicable.

Note: After the completion of this work, Refs. [17, 18] were brought to my attention where the authors use two different two-component formulations of the Wheeler-DeWitt equation. The methods described in these articles differ substantially from the present article's. I have also noticed a recent paper by Embacher [19] who defines the operator D of Eq. (7) as the (matter) energy operator and obtains the mode expansion (32) in the context of the one-component Wheeler-DeWitt equation. He does not, however, obtain the equations satisfied by the mode

coefficients. Instead, he focuses on the semiclassical and adiabatic approximations. The two-component approach developed in this paper justifies the choice of the mode functions made by Embacher in a very natural way. It also makes it possible to obtain the structure of the mode expansion by iteratively solving for the mode coefficients.

Acknowledgements

I would like to thank B. Darian, M. Razavi, and especially D. Page for invaluable comments and suggestions. I would also like to acknowledge the financial support of the Killam Foundation of Canada.

References

- [1] D. N. Page, “Lectures on Quantum Cosmology,” in *Gravitation: A Banff Summer Institute*, edited by R. Mann and P. Wesson (World Scientific, Singapore, 1991);
D. L. Wiltshire, “An Introduction to Quantum Cosmology,” in *Cosmology: The Physics of the Universe*, Adelaide preprint no ADP-95-11/M28 (World Scientific, Singapore, to appear).
- [2] D. J. Kaup and A. P. Vitello, Phys. Rev. **D 9**, 1648 (1974).
- [3] S. W. Hawking and D. N. Page, Phys. Rev. **D42**, 2655, (1990).
- [4] A. Mostafazadeh, “Adiabatic Geometrical Phase for Scalar Fields in a Curved Spacetime,” University of Alberta preprint no: THY-29-96, hep-th/9608051 (1996).
- [5] H. Feshbach and F. Villars, Rev. Mod. Phys. **30**, 24 (1958);
B. R. Holstein, *Topics in Advanced Quantum Mechanics* (Addison-Wesley, Redwood City, California, 1992).
- [6] R. Jackiw, Int. J. Mod. Phys. **A 3**, 285 (1988).
- [7] L. I. Schiff, *Quantum Mechanics* (McGraw-Hill, New York, 1955).

- [8] A. Mostafazadeh, “The Quantum Adiabatic Approximation and the Geometric Phase,” Phys. Rev. **A**, to appear, University of Alberta preprint no: THY-25-96, hep-th/9606053.
- [9] A. Mostafazadeh, “Adiabatic Product Expansion,” University of Alberta preprint no: THY-26-96, quant-ph/9606032 (1996).
- [10] R. M. Wald, *Quantum Field Theory in Curved Spacetime and Black Hole Thermodynamics*, Appendix A.1 (University of Chicago Press, Chicago, 1994).
- [11] C. Kiefer, Phys. Rev. **D 38**, 1761 (1988).
- [12] J. B. Hartle and S. W. Hawking, Phys. Rev. **D28**, 2960 (1983).
- [13] I. S. Gradshteyn and I. M. Ryzhik, *Table of Integrals, Series, and Products*, (Academic Press, New York, 1980).
- [14] T. Banks, Nucl. Phys. **B249**, 332 (1985);
R. Brout, Found. Phys. **17**, 603 (1987);
R. Brout and G. Venturi, Phys. Rev. **D39**, 2436 (1989);
G. Venturi, “Quantum Gravity and the Berry Phase,” in *Differential Geometric Methods in Theoretical Physics*, Eds. L.-L. Chau and W. Nahm (Plenum, New York, 1990);
D. P. Datta, Phys. Rev. **D 48**, 5746 (1993).
- [15] B. S. DeWitt, Phys. Rev. **160**, 1113 (1967).
- [16] H. R. Lewis, Jr. and W. B. Riesenfeld, J. Math. Phys. **10**, 1458 (1969);
D. M. Greenberger, J. Math. Phys. **15**, 395 (1974);
R. W. Hasse, J. Math. Phys. **16**, 2005 (1975);
V. V. Dodonov and V. I. Man’ko, Phys. Rev. **A 20**, 550 (1979);
L. F. Landovitz, A. M. Levine and W. M. Schreiber, Phys. Rev. **A 20**, 1162 (1979);
M. S. Abdalla and R. K. Colegrave, Phys. Rev. **A 32**, 1958 (1985);
A. B. Nassar and R. T. Berg, Phys. Rev. **A 34**, 2462 (1986);
C. M. Cheng and P. C. W. Fung, J. Phys. A:Math. Gen. **21**, 4115 (1988);
F. Wolfs and H. J. Korsch, Phys. Rev. **A 37**, 1934 (1988);

- G. S. Agarwal and S. Arun Kumar, Phys. Rev. Lett. **67**, 3665 (1991);
- A. de Souza Dutra, C. F. de Souza, and L. C. de Albuquerque, Phys. Lett. **A 156**, 371 (1991);
- S. P. Kim, J. Phys. A: Math. Gen. **27**, 3927 (1994);
- J.-Y. Ji and J. K. Kim, Phys. Rev. **A 51**, 4268 (1995).
- [17] S. P. Kim, Phys. Rev. **D 46**, 3403 (1992).
- [18] T. Dereli, M. Önder, and R. W. Tucker, Phys. Lett. **B324**, 134 (1994).
- [19] Franz Embacher, “Quantum Cosmology in the Energy Representation,” Uni. Wien preprint no: UWThPh-1996-32, gr-qc/9605021.




Figure 1: Plot of $\psi_0 = \psi_2 = \sum_{n=0}^2 \sigma_n \Phi_n$, for $\sigma_{\pm 1} = \sigma_{-0} = 0$, $\sigma_0 = 1$ and $m = 10^{-6}$.




Figure 2: Plot of $\psi_4 := \sum_{n=0}^4 \sigma_n \Phi_n$, for $\sigma_{\pm 1} = \sigma_{-0} = 0$, $\sigma_0 = 1$ and $m = 10^{-6}$.




Figure 3: Plot of $\psi_6 := \sum_{n=0}^6 \sigma_n \Phi_n$, for $\sigma_{\pm 1} = \sigma_{-0} = 0$, $\sigma_0 = 1$ and $m = 10^{-6}$.




Figure 4: Plot of $\psi_8 := \sum_{n=0}^8 \sigma_n \Phi_n$, for $\sigma_{\pm 1} = \sigma_{-0} = 0$, $\sigma_0 = 1$ and $m = 10^{-6}$.




Figure 5: Plot of $\psi_{10} := \sum_{n=0}^{10} \sigma_n \Phi_n$, for $\sigma_{\pm 1} = \sigma_{-0} = 0$, $\sigma_0 = 1$ and $m = 10^{-6}$.




Figure 6: Plot of the restriction of ψ_6 (dotted line), ψ_8 (dashed line) and ψ_{10} (full line) to the $a = 0.1$ line. Note that the graphs of ψ_8 and ψ_{10} coincide for this value of a . This shows the reliability of the finite mode approximation. Note also the range of the values of ϕ over which ψ_n are non-zero.




Figure 7: Plot of the restriction of ψ_6 (dotted line), ψ_8 (dashed line) and ψ_{10} (full line) to the $a = 1$ line. Note the range of the values of ϕ over which ψ_n are non-zero.




Figure 8: Plot of the restriction of ψ_8 (dotted line), and ψ_{10} (full line) to the $a = 10$ line. Note the range of the values of ϕ over which ψ_n are non-zero.




Figure 9: Plot of the restriction of ψ_8 (dotted line), and ψ_{10} (full line) to the $a = 100$ line. Note the range of the values of ϕ over which ψ_n are non-zero.